

R15

Code No: 121AL

JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY HYDERABAD

B. Tech I Year Examinations, January/February - 2024

MATHEMATICAL METHODS

(Common to EEE, ECE, CSE, EIE, IT)

Time: 3 Hours

Max. Marks: 75

Note: i) Question paper consists of Part A, Part B.

ii) Part A is compulsory, which carries 25 marks. In Part A, answer all questions.

iii) In Part B, Answer any one question from each unit. Each question carries 10 marks and may have a, b as sub questions.

PART - A

(25 Marks)

- 1.a) Write the normal equations to fit the parabola $y = a + bx + cx^2$ using least squares method. [2]
- b) Determine $(2\Delta + 3)(E + 2)(3x^2 + 2)$ by taking $h = 1$. [3]
- c) Explain Trapezoidal to find the approximate value of $\int_a^b f(x)dx$. [2]
- d) Derive the recurrence relation to evaluate \sqrt{N} using Newton Raphson's method. [3]
- e) Is $f(x) = \sin x + \frac{1}{3}\sin 3x + \frac{1}{5}\sin 5x + \frac{1}{7}\sin 7x$ periodic? If so, find its period. [2]
- f) Obtain the Fourier sine integral of $f(x) = e^{-kx}$, $k > 0$ & $x \geq 0$. [3]
- g) Using the method of separation of variables, find solution of $x^2 \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 0$. [2]
- h) Form a partial differential equation by eliminating arbitrary constants a, b from $2z = \sqrt{(x+a)} + \sqrt{(y-a)} + b$. [3]
- i) For what value(s) of p , the vector field $\vec{f} = (x+3y)\vec{i} + (y-2z)\vec{j} + (x+pz)\vec{k}$ is solenoid vector? [2]
- j) Compute the work done in moving a particle in the force field $\vec{F} = 3x^2\vec{i} + \vec{j} + z\vec{k}$ along the straight line from $(0,0,0)$ to $(2,1,3)$. [3]

PART - B

(50 Marks)

- 2.a) Prove or disprove: $\mu^2 = 1 + \frac{\delta^2}{4}$.
- b) Use Lagrange's interpolation formula to fit a polynomial to the following data.

x	-1	0	2	3
y	-8	3	1	2

Hence find $y(-2)$, $y(1)$ and $y(4)$.

[4+6]

OR

- 3.a) Explain the method of least squares.
 b) Determine the constants a and b by the method of least squares such that $y = ae^{bx}$.

x	2	4	6	8	10
y	4.077	11.084	30.128	81.897	222.62

- 4.a) Compute a positive real root of the equation $x \log_{10} x = 1.2$ correct to three decimal places using Bisection method.

- b) Using the Gauss Seidel iterative method, solve:

$$5x - y = 9; x - 5y + z = -4; y - 5z = 6. \quad [5+5]$$

5. Use a) Taylor series method b) Runge-Kutta method of 4th order to solve

$$10 \frac{dy}{dx} = x^2 + y^2, y(0) = 1 \text{ for the interval } 0 < x \leq 0.2 \text{ with } h = 0.1. \quad [5+5]$$

6. Find the Fourier series of periodic function $f(x)$ of period 2, where

$$f(x) = \begin{cases} -1 & \text{if } -1 < x < 0 \\ 2x & \text{if } 0 \leq x < 1 \end{cases} \quad \text{Hence, prove that } 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots = \frac{\pi}{4}. \quad [10]$$

- 7.a) State the Fourier integral theorem.

- b) Represent $f(x)$ as an exponential Fourier transform when $f(x) = \begin{cases} \sin x, & 0 < x < \pi \\ 0, & \text{otherwise} \end{cases}$.

$$\text{Show that the result can be written as } f(x) = \frac{1}{\pi} \int_0^{\infty} \frac{\cos \lambda x - \cos \lambda(x - \pi)}{1 - \lambda^2} d\lambda. \quad [2+8]$$

8. Solve the following differential equations

a) $(z^2 - 2yz - y^2)p + (xy + xz)q = xy - xz$, where $p = \frac{\partial z}{\partial x}, q = \frac{\partial z}{\partial y}$.

b) $(p^2 + q^2)y = qz$, where $p = \frac{\partial z}{\partial x}, q = \frac{\partial z}{\partial y}$. [4+6]

9. A string is stretched and fastened to two points l apart. Motion is started by displacing the string in the form $y = a \sin(\pi x/l)$ from which it is released at time $t = 0$. Show that the displacement of any point at a distance x from one end at time t is given by

$$y(x, t) = a \sin\left(\frac{\pi x}{l}\right) \cos\left(\frac{\pi ct}{l}\right) \quad [10]$$

- 10.a) Calculate the scalar potential ϕ such that $\vec{f} = \nabla \phi$ if the vector field $\vec{f} = (x^2 - yz)\vec{i} + (y^2 - zx)\vec{j} + (z^2 - xy)\vec{k}$ is an irrotational vector.

- b) Evaluate the directional derivative of $f = x^2 - y^2 + 2z^2$ at the point P(1, 2, 3) in the direction of the line PQ where Q is the point (5, 0, 4). [5+5]

11. Verify the divergence theorem for $\vec{f} = (x^2 - yz)\vec{i} + (y^2 - zx)\vec{j} + (z^2 - xy)\vec{k}$ taken over the rectangular parallelepiped $0 \leq x \leq a, 0 \leq y \leq b, 0 \leq z \leq c$. [10]